

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C. 20231
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

| | |
|---|---|
| Date of mailing (day/month/year) 24 February 2000 (24.02.00) | |
| International application No. PCT/GB99/00905 | Applicant's or agent's file reference KVC/P20895WO |
| International filing date (day/month/year) 22 March 1999 (22.03.99) | Priority date (day/month/year) 20 March 1998 (20.03.98) |
| Applicant WYATT, Paul et al | |

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

19 October 1999 (19.10.99)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

| | |
|--|---|
| The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland | Authorized officer Maria Victoria CORTIELLO |
| Facsimile No.: (41-22) 740.14.35 | Telephone No.: (41-22) 338.83.38 |

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

| | | |
|--|---|--|
| Applicant's or agent's file reference KVC/P20895WO | <div style="display: flex; justify-content: space-between;"> <div>FOR FURTHER ACTION</div> <div>See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)</div> </div> | |
| International application No. PCT/GB99/00905 | International filing date (<i>day/month/year</i>) 22/03/1999 | Priority date (<i>day/month/year</i>) 20/03/1998 |
| International Patent Classification (IPC) or national classification and IPC C12N15/29 | | |
| Applicant BIOGEMMA UK LIMITED et al. | | |

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 7 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

| | |
|---|--|
| Date of submission of the demand 19/10/1999 | Date of completion of this report 16.06.2000 |
| Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465 | Authorized officer Burkhardt, P Telephone No. +49 89 2399 7456  |

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/GB99/00905

I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

Description, pages:

1-36 as originally filed

Claims, No.:

1-29 as originally filed

Drawings, sheets:

1/17-17/17 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

see separate sheet

II. Priority

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:
- ☐ copy of the earlier application whose priority has been claimed.
 - ☐ translation of the earlier application whose priority has been claimed.
2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/GB99/00905

been found invalid.

Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:

see separate sheet

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

| | | | |
|-------------------------------|------|--------|---|
| Novelty (N) | Yes: | Claims | 4, 5, 9, 12, 14 - 16, 18, 20 - 24 |
| | No: | Claims | 1 - 3, 6 - 8, 10, 11, 13, 17, 19, 25 - 29 |
| Inventive step (IS) | Yes: | Claims | 4, 5, 18, 20 - 24 |
| | No: | Claims | 9, 12, 14 - 16 |
| Industrial applicability (IA) | Yes: | Claims | 1 - 29 |
| | No: | Claims | |

2. Citations and explanations

see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:

see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

see separate sheet

Re Item I

Basis of the opinion

This written report is also based on the Sequence Listing (pages 1-25) as filed with the letter of 01.07.1999.

Re Item II

Priority

As the priority document was not available at the time of drafting the present written opinion, this opinion is being established under the assumption that the entire subject-matter is entitled to the claimed priority. If this is not the case, the 'P,X' document cited in the search report may have to be considered for novelty and/or inventive step. This document would be detrimental to most, if not all, of the claimed subject-matter (Article 33(2), (3) PCT).

Re Item V

Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

The following document (D) is referred to in this report; the numbering is following the order of the International Search Report:

D1 Nelson *et al.*, 1997. Plant Physiol. 114:29-37.

1. Subject-matter of the application

The applicant has identified and cloned the gene DZ2 from *Brassica napus* and its orthologue from *Arabidopsis thaliana*. The cDNA was isolated by differential display from pod dehiscence zone tissue. A set of claims based on DZ2 from *Arabidopsis* (SEQ ID NO 29) and *Brassica* (SEQ ID NO 23) appears to be novel and inventive and would therefore meet the requirements of Article 33(2)(3) PCT. The claims in their present form, however, give rise to the following objections. For the interpretation of the claims please also see section VIII 1.

2. Article 33(2)(3) PCT (Novelty and inventive step)

2.1 Present claim 1 is directed to a nucleic acid encoding a signal transduction protein involved in the process of dehiscence.

Prior art document D1 discloses nucleic acid sequences (page 32, Table 3) that may be involved in dehiscence (page 29, abstract) and which encodes a signal transduction protein (page 29, second paragraph). D1 furthermore discloses the expression of these genes in *E. coli*, the isolation of the corresponding proteins and antibodies against these proteins.

Thus, D1 anticipates the novelty of the subject-matter of present claim 1. The same holds true for dependent claims 2, 3, 6 - 8, 10 and 11. D1 also anticipates the subject-matter of present claims 13, 17 and 19 addressing a process for obtaining a cell containing said nucleic acid and a signal transduction protein as well as present claims 25, 26 and 29 directed to the use of said nucleic acids or proteins as probes or for the production of a cell.

2.2 Present claim 27 relates to a nucleic acid sequence comprising one or more of the underlined sequences as set out in Figure 1. These sequences include *NcoI* and *HindIII* restriction sites that are present in numerous nucleotide sequences. Many of these sequences definitely have been used as PCR primers. Thus, the subject-matter of present claims 27 and 28 is not novel over the prior art.

2.3 Present claims 4 and 5 are directed to nucleic acid sequences encoding a protein that comprises the amino acid sequence shown in Figure 1, or modifications or fragments thereof. Such sequences have not yet been disclosed nor suggested in the prior art presently available to the IPEA. The same holds true for the protein itself as set out in present claim 18

2.4 Present claim 9 is directed to the regulator/promoter sequence of a nucleic acid sequence according to claims 1 - 8. Such a sequence has not yet been disclosed in the prior art presently available to the IPEA.

The nucleic acid sequences of claims 1 - 3, however, are not novel. It cannot be regarded as involving an inventive step to isolate the promoter of a known nucleic acid sequence. It is within the scope of an ordinary man skilled in the art to isolate and characterise genomic sequences once cDNA clones are available

(see 2.1).

2.5 A similar objection applies to present claim 12. It cannot be regarded as involving an inventive step to transform plant cells with known sequences. Methods for plant transformation are readily available and can be used to transfer any known nucleic acid sequence. Thus, claim 12 does not meet the requirements of Article 33(3) PCT. The same holds true for present claims 14 - 16.

2.6 Present claim 20 is directed to a process of regulating/controlling dehiscence in a plant or a part thereof by transforming the plant with nucleic acid sequences as set out in claims 1 - 10. Such a process has not yet been disclosed nor suggested in the prior art presently available to the IPEA. Thus, claim 20 meets the requirements of Article 33(2)(3) PCT. The same holds true for present claims 21 - 24.

Re Item VII

Certain defects in the international application

1. The reference to claim 21 within present claim 21 appears to be incorrect.

Re Item VIII

Certain observations on the international application

1. The applicants have isolated and sequenced *Arabidopsis* and *Brassica* genes which are homologous to bacterial response regulator proteins. However, the analysis of the postulated function, i.e. encoding the receiver domain of a response regulator, is not provided. Consequently, claims relating to nucleic acids and polypeptides involved in dehiscence, i.e. present claims 1 - 3 and claims referring to these claims, are not supported by the description (Article 6 PCT). It appears to be furthermore impossible for a person skilled in the art to arrive at the claimed subject-matter without undue experimentation (Article 5 PCT).
2. Based on the cloning and sequencing of DZ2 it appears to be unjustified to extend

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/GB99/00905

the scope of the claims to structurally unrelated proteins equally involved in dehiscence. The applicants did not contribute to the cloning of genes other than DZ2.

A similar objection applies to present claim 6 addressing combinations with further nucleic acid sequences that are expressed in the dehiscence zone.

3. The term "involved" in present claims 1 and 17 is unclear (Article 6 PCT). It could be interpreted as embracing various cellular housekeeping proteins that all would anticipate the subject-matter of present claims 1 and 17.




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|--|--|--|---|
| (51) International Patent Classification ⁶ : C12N 15/29, 15/82, 15/11, 5/10, C07K 14/415, A01H 5/00, 5/10 | | A1 | (11) International Publication Number: WO 99/49046 |
| (21) International Application Number: PCT/GB99/00905 | | (43) International Publication Date: 30 September 1999 (30.09.99) | |
| (22) International Filing Date: 22 March 1999 (22.03.99) | | | |
| (30) Priority Data: 9806113.8 20 March 1998 (20.03.98) GB | | | |
| (71) Applicant (for all designated States except US): BIOGEMMA UK LIMITED [GB/GB]; 200 Science Park, Milton Road, Cambridge CB4 0GZ (GB). | | (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). | |
| (72) Inventors; and (75) Inventors/Applicants (for US only): WYATT, Paul [GB/GB]; Biogemma UK Limited, 200 Science Park, Milton Road, Cambridge CB4 0GZ (GB). ROBERTS, Jeremy, Alan [GB/GB]; University of Nottingham, Division of Plant Science, School of Biological Sciences, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD (GB). WHITELAW, Catherine [GB/GB]; University of Nottingham, Division of Plant Science, School Of Biological Sciences, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD (GB). | | Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. | |
| (74) Agents: CORNISH, Kristina, Victoria, Joy et al.; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB). | | | |

(54) Title: SIGNAL TRANSDUCTION PROTEIN INVOLVED IN PLANT DEHISCENCE

Expression analysis of DZ2 in various plant organs using Northern



(57) Abstract

This invention relates to novel plant nucleic acid sequences and proteins. The sequences and proteins are useful in the control of plant dehiscence and in the production of male sterile plants. According to a first aspect of the invention there is provided nucleic acid optionally encoding a signal transduction protein involved in the process of dehiscence. Such a sequence or signal transduction protein has never previously been described in plant dehiscence.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

**SIGNAL TRANSDUCTION PROTEIN INVOLVED
IN PLANT DEHISCENCE**

5 This invention relates to novel plant nucleic acid sequences and proteins. The sequences and proteins are useful in the control of plant dehiscence and in the production of male sterile plants.

10 The production of seed is an important developmental process in all higher plants. In oilseed rape (*Brassica napus*), following abscission of floral parts, pods or siliques are formed which contain 15-30 seeds. Around 50-70 days after anthesis (DAA) the pods become susceptible to shatter, a process that serves to expel the mature seeds into the surrounding environment. In the days leading to dehiscence, an array of anatomical, molecular and biochemical changes take place, thus preparing both seed and pod for the event. Shatter eventually occurs as a result of a
15 combination of factors including: the creation of tensions within the pod between the lignified valve edge cells of the endocarp and the unlignified dehiscence zone (DZ) cells, weakening of the DZ cell walls by hydrolytic enzyme activity and ultimately due to physical forces such as strong winds or harvesting machinery.

20 Pod development in *B. napus* can be segmented into three stages. In the first stage, which occurs 0-20 DAA, the newly formed siliques, consisting of two seed-containing carpels separated by a false septum and a replar region, grow to their full length of around 10cm. The seeds begin to grow when the pods are virtually full size [Hocking and Mason, 1993]. Between 10 and 20 DAA the cells in the replar
25 region begin to differentiate into replar cells, large valve edge cells and form a distinct region, 1-3 cells wide, comprising the DZ [Meakin and Roberts, 1990a].

The second stage occurs between 20 and 50 DAA. From 20 DAA, in conjunction

- with termination of pod elongation, secondary cell wall material is deposited in the valve edge cells, and the replar cells become increasingly lignified. The DZ cells do not exhibit thickening of the cell wall. A progressive shrinkage and loss of organelles is apparent in the DZ cells from 40 DAA onwards and eventually these cells separate completely due to hydrolysis of the middle lamella [Meakin and Roberts, 1990a]. In the final stage of pod development, which occurs 50-70 DAA, the lignified cells undergo senescence and the necessary tensions are created so that the desiccated pod, containing mature seed, eventually shatters.
- 10 Molecular studies of the penultimate stage of pod development have revealed a spatial and temporal correlation between the up-regulation of a number of mRNAs and pod dehiscence in *B. napus*. These mRNAs encode a polygalacturonase (PG) and a proline-rich protein (SAC51). Further analysis of the expression of the PG following fusion of a pod-specific *Arabidopsis thaliana* PG promoter to GUS
- 15 [Jenkins et al., (1997)], has revealed that reporter gene expression is restricted precisely to the layer of cells comprising the pod DZ in transgenic *B. napus*. From 40 DAA, Meakin and Roberts (1990b) reported a progressive increase in β -1,4-glucanase (cellulase) activity in the DZ.
- 20 It is understood that the processes of dehiscence and abscission are not regulated by the same environmental or chemical signals, but that they involve controlled degradation of cell wall material and cell separation in a distinct group of cells. Both ethylene and indole-3-acetic acid (IAA) appear to be important regulators of the timing of the abscission process but the role of these plant hormones in dehiscence is
- 25 less clearly defined. The increase in cellulase activity has been shown to correlate with a rise in the production of ethylene, mainly from the seed, which peaks at around 40 DAA [Meakin and Roberts, 1990b; Johnson-Flanagan and Spencer, 1994].

Developmental processes, such as pod dehiscence, which involve highly regulated and controlled expression of an array of different genes at a precise time and cellular location, clearly require an intricate signal transduction network.

5 Further and improved genetic elements to control plant processes in this area are constantly desired. We describe the isolation, for the first time, of a plant cDNA (DZ2) encoding an individual response regulator protein, the expression of which is closely correlated with dehiscence of fruit in *B. napus*. DZ2 has a role in the ability to control molecule signaling during the events leading to shatter and thus to control
10 pod shatter in plants. In addition to the identification of the nucleic acid termed "DZ2" a homologous, but not identical sequence and protein were also identified from *B. napus*. This sequence was designated "DZ2B". Sequence analysis of DZ2 and DZ2B shows that there are two DZ2 genes in *B. napus*, each represented by a slightly different cDNA (here termed DZ2 and DZ2B). This is consistent with one
15 gene being encoded by the *B. campestris* derived genome and the other from the genome derived from *B. oleracea*. In this text, the designation "DZ2" is equivalent to the CW1 designation in UK 9806113.8 (as seen from Figure 1).

According to a first aspect of the invention there is provided nucleic acid optionally
20 encoding a signal transduction protein involved in the process of dehiscence. Such a sequence or signal transduction protein has never previously been described in plant dehiscence.

In this text, the term "involved in the process of dehiscence" means any nucleic acid
25 (preferably) encoding any protein which has an effect in the dehiscence process, in particular a protein or nucleic acid sequence involved in an MAP Kinase cascade or any other protein or nucleic acid sequence which results in changes in the expression of genes involved in dehiscence, such as upregulation of genes encoding

polygalacturonase, cellulase, senescence-related proteins and/or downregulation of genes encoding for proteins involved in cell wall biosynthesis. The nucleic acid sequences/proteins of the present invention which are "involved in the process of plant dehiscence" are not the individual structural genes or proteins which cause dehiscence (polygalacturonases etc.). Rather, the nucleic acid sequences/proteins of the present invention are sequences/proteins which have an effect on the expression of such structural genes or proteins. One advantage of the present invention is that the use of such nucleic acid sequences/proteins enables the possibility to influence the whole process of dehiscence rather than just one element of it. Preferably the protein or nucleic acid sequence of the present invention which is involved in the process of dehiscence effects a structural protein which is a hydrolytic enzyme such as polygalacturonase or cellulase.

The nucleic acid of the first aspect of the invention may be a nucleic acid which is naturally expressed in a dehiscence zone. Such a nucleic acid will most accurately reflect nucleic acid naturally expressed in a plant. Preferably the dehiscence zone is a pod (also termed "siliques"), anther and/or funiculus dehiscence zone. Preferably the plant is a member of the Brassica family, maize, wheat, soyabean, *Cuphea* or sesame.

20

A second aspect of the invention provides nucleic acid encoding a protein wherein the protein:

- a) comprises the amino acid sequence shown in figure 1 or;
- b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above, but has at least 40% amino acid sequence identical therewith; or

25

- c) is a fragment of a protein as defined in a) or b) above, which is at least 10 (preferably 20 or 30) amino acids long.

5 The percentage amino acid identity can be determined using the default parameters of the GAP computer program, version 6.0 described by Deveraux *et al.*, 1984 and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilises the alignment method of Needleman and Wunsch 1970 as revised by Smith and Waterman 1981. More preferably the protein has at least 45%
10 identity to the amino acid sequence of Figure 1, through 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% identity using the default parameters.

The skilled person will appreciate that various changes can sometimes be made to the amino acid sequence of a protein (which has a desired property) to produce
15 variants (often known as "muteins") which still have said property. Such variants of the protein describe in a, b and c above are within the scope of the present invention and are discussed in greater detail below in sections (i) to (iii). They include allelic and non-allelic variants.

20 (i) *Substitutions*

An example of a variant of the present invention is a polypeptide as defined in a, b or c above, apart from the substitution of one or more amino acids with one or more other amino acids.

25 The skilled person is aware that various amino acids have similar characteristics. One or more such amino acids of a protein can often be substituted by one or more other such amino acids without eliminating a desired property of that protein.

For example, the amino acids glycine, alanine, valine, leucine and isoleucine can often be substituted for one another (amino acids having aliphatic side chains). Of these possible substitutions it is preferred that glycine and alanine are used to substitute for one another (since they have relatively short side chains) and that
5 valine, leucine and isoleucine are used to substitute for one another (since they have larger aliphatic side chains which are hydrophobic). Other amino acids that can often be substituted for one another include phenylalanine, tyrosine and tryptophan (amino acids having aromatic side chains); lysine, arginine and histidine (amino acids having basic side chains); aspartate and glutamate (amino acids having acidic
10 side chains); asparagine and glutamine (amino acids having amide side chains); and cysteine and methionine (amino acids having sulphur containing side chains).

Substitutions of this nature are often referred to as "conservative" or "semi-conservative" amino acid substitutions.

15

(ii) *Deletions*

Amino acid deletions can be advantageous since the overall length and the molecular weight of a polypeptide can be reduced whilst still retaining a desired property. This can enable the amount of protein required for a particular purpose to be reduced.
20 Proteins according to the present invention, which have such deletion(s) are useful. They may interfere with the normal functioning of DZ2; that is, they may act as dominant negative mutations preventing normal DZ2 functioning and thus be of particular value, for example, in reducing pod shatter.

25 The amino acid sequence shown in figure 1 has various functional regions. For particular applications of the present invention, one or more of these regions may not be needed and may therefore be deleted.

(iii) *Insertions*

Amino acid insertions relative to a polypeptide as defined in a, b or c above can also be made. This may be done to alter the nature of the protein (e.g. to assist in identification, purification, or expression, as explained below in relation to fusion proteins).

Changes in the protein according to the present invention can produce versions of the protein that are constitutively active. If a protein of the present invention acts on an inhibitor of the release of hydrolytic enzymes, then a constitutively active version would prevent or reduce pod shatter

A protein according to any aspect of the invention may have additional N-terminal and/or C-terminal amino acid sequences. Such sequences can be provided for various reasons. Techniques for providing such sequences are well known in the art. They include using gene-cloning techniques to ligate together nucleic acid molecules encoding polypeptides or parts thereof, followed by expressing a polypeptide encoded by the nucleic acid molecule produced by ligation.

Additional sequences may be provided in order to alter the characteristics of a particular polypeptide. This can be useful in improving expression or regulation of expression in particular expression systems. For example, an additional sequence may provide some protection against proteolytic cleavage. This has been done for the hormone somatostatin by fusing it at its N-terminus to part of the β galactosidase enzyme [Itakwa *et al.*, 105-63 (1977)].

Additional sequences can also be useful in altering the properties of a polypeptide to aid in identification or purification.

For example, a signal sequence may be present to direct the transport of the polypeptide to a particular location within a cell or to export the polypeptide from the cell. Hydrophobic sequences may be provided to anchor a polypeptide in a membrane. Thus the present invention includes within its scope both soluble and
5 membrane-bound polypeptides.

Preferably, the nucleic acid according to the second aspect of the invention encodes a signal transduction protein or a functional portion thereof involved in the process of dehiscence. All preferred features of the first aspect of the invention as described
10 above also apply to the second.

The term protein used in this text means, in general terms, a plurality of amino acid residues joined together by peptide bonds. It is used interchangeably and means the same as polypeptide or peptide.

15

The nucleic acid according to the first or second aspect of the invention preferably comprises the sequence set out in figure 1 or a sequence which is 40% or more identical, preferably through 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% to the sequence in Figure 1 at the nucleic acid residue level, using the
20 default parameters of the GAP computer program, version 6.0 described by Deveraux *et al.*, 1984 and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilises the alignment method of Needleman and Wunsch 1970 as revised by Smith and Waterman 1981. Further, the nucleic acid may comprise a fragment of a sequence according to the first or second
25 aspect which is at least 30 bases long also 40, 50, 60, 70, 80, or 90 bases in length. While this nucleic acid is the preferred nucleic acid of the invention, it is well known to those persons skilled in the art that because of the nucleic acid "degenerate code" which encodes nucleic acids, a considerable number of variations in nucleic acid

sequence can be used to encode for proteins according to the first or second aspects of the invention.

5 The nucleic acid of the first or second aspects of the invention may be isolated or recombinant and may be in substantially pure form. The nucleic acid may be antisense to nucleic acid according to the first or second aspects of the invention. As understood by the person skilled in the art introducing the coding region of a gene in the reverse orientation to that found in nature (antisense) can result in the downregulation of the gene and hence the production of less or none of the gene product. The transcribed antisense DNA is capable of binding to and destroying the function of the sense RNA of the sequence normally found in the cell, thereby disrupting function. Antisense nucleic acid may be constitutively expressed, but is preferably limited to expression in those zones (dehiscence) in which the naturally occurring nucleic acid is expressed.

15 The nucleic acid according to the first or second aspects of the invention preferably include a promoter or other regulatory sequence which controls expression of the nucleic acid. Promoters and other regulatory sequences which control expression of a nucleic acid in dehiscence zones are known in the art, for example described in WO96/30529 and WO94/23043. Further promoters or other regulatory sequences can be identified and can also include the promoter or other regulatory sequence which controls expression of a nucleic acid as set out in figure 1. The person skilled in the art will know that it may not be necessary to utilize the whole promoter or other regulatory sequence. Only the minimum essential regulatory elements may be required and in fact such elements can be used to construct chimeric sequences or promoters. The essential requirement is, of course, to retain the tissue and/or temporal specificity.

The nucleic acid according to the first or second aspects of the invention may be in the form of a vector. The vector may be a plasmid, cosmid or phage. Vectors frequently include one or more expressed markers which enable selection of cells transfected (or transformed: the terms are used interchangeably in this text) with them and preferably, to enable a selection of cells containing vectors incorporating heterologous DNA. A suitable start and stop signal will generally be present and if the vector is intended for expression, sufficient regulatory sequences to drive expression will be present. Nucleic acid according to the first and second aspects of the invention is preferably for expression in plant cells and thus microbial host expression is perhaps less important although not ruled out. Microbial host expression and vectors not including regulatory sequences are useful as cloning vectors.

A third aspect of the invention relates to a cell comprising nucleic acid according to the first or second aspects of the invention. The cell may be termed as "a host" which is useful for manipulation of the nucleic acid, including cloning. Alternatively, the cell may be a cell in which to obtain expression of the nucleic acid, most preferably a plant cell. The nucleic acid can be incorporated by standard techniques known in the art in to cells. Preferably nucleic acid is transformed in to plant cells using a disarmed Ti plasmid vector and carried by an Agrobacterium by procedures known in the art, for example as described in EP-A-0116718 and EP-A-0270822. Foreign nucleic acid can alternatively be introduced directly into plant cells using an electrical discharged apparatus or by any other method that provides for the stable incorporation of the nucleic acid into the cell. Preferably the stable incorporation of the nucleic acid is within the nucleic DNA of any cell preferably a plant cell. Nucleic acid according to the first and second aspects of the invention preferably contains a second "marker" gene that enables identification of the nucleic acid. This is most commonly used to distinguish the transformed plant cell

- containing the foreign nucleic acid from other plants cells that do not contain the foreign nucleic acid. Examples of such marker genes include antibiotic resistance, herbicide resistance and Glucuronidase (GUS) expression. Expression of the marker gene is preferably controlled by a second promoter which allows expression of the
- 5 marker gene in cells other than those than dehiscence zones (if this is the tissue specific expression of the nucleic acid according to the first or second aspects of the invention). Preferably the cell is from any of the Brassica family (most preferably *B. napus*), maize, wheat, soyabean, *Cuphea* and sesame.
- 10 A third aspect of the invention includes a process for obtaining a cell comprising nucleic acid according to the first or second aspects of the invention. The process involves introducing said nucleic acid into a suitable cell and optionally growing on or culturing said cell.
- 15 A fourth aspect of the invention provides a plant or a part thereof comprising a cell according to the third aspect of the invention. A whole plant can be regenerated from the single transformed plant cell by procedures well known in the art. The invention also provides for propagating material or a seed comprising a cell
- 20 according to the third aspect of the invention. The invention also relates to any plant or part thereof including propagating material or a seed derived from any aspect of the invention. The fourth aspect of the invention also includes a process for obtaining a plant or plant part (including propagating material or seed, the process comprising obtaining a cell according to the third aspect of the invention or, indeed, plant material according to the fourth aspect of the invention and growth (to the
- 25 required plant, plant part, propagating material etc). Techniques for such a process are commonplace in the art.
- A fifth aspect of the invention provides a signal transduction protein involved in the

process of the plant dehiscence. The signal transduction protein according to the fifth aspect may have one or more of the preferred features according to the first or second aspects of the invention. Preferably it may be isolated, recombinant or in substantially pure form. It may comprise the various changes according to the first or second aspects. Preferably the protein is expressed from nucleic acid according to the first or second aspects. Alternatively, the protein can be provided using suitable techniques known in the art.

A sixth aspect of the invention provides a protein which;

- a) comprises the amino acid sequence shown in figure 1 or;
- b) has one or more amino acid deletions, insertions, or substitutions relative to a protein as defined in a) above and has at least 40% amino acid sequence identity therewith;

or a fragment of a protein as defined in a) or b) above which is at least 10 amino acids long. The percentage amino acid identity can be determined using the default parameters of the GAP computer program, version 6.0 described by Deveraux *et al.*, 1984 and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilises the alignment method of Needleman and Wunsch 1970 as revised by Smith and Waterman 1981. More preferably the protein has at least 45% identity to the amino acid sequence of Figure 1, through 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% identity using the default parameters.

The protein is preferably a signal transduction protein involved in the process of plant dehiscence and again, the preferred features of aspects one, two and five also applied to the sixth aspect.

The seventh aspect of the invention provides a process for regulating/controlling dehiscence in plant or in a part thereof, the process comprising obtaining a plant or a part thereof according to the fourth aspect of the invention. The process of dehiscence can be regulated and/or controlled by increasing or decreasing the expression of nucleic acid sequences according to the first or second aspect of the invention. Increased or decreased expression can easily be influenced by the person skilled in the art using technology well known. This includes increasing the numbers of copies of nucleic acid according to the invention in a plant or a plant thereof or increasing expression levels of copies of the nucleic acid present in particular parts or zones of the plant. Preferably the zones are dehiscence zones.

The process according to the seventh aspect of the invention includes obtaining a plant cell according to the third aspect of the invention or part of a plant according to the fourth aspect in the invention and deriving a plant therefrom. Alternatively, the process may comprise obtaining propagating material or a seed according to the fourth aspect of the invention and deriving a plant therefrom.

Preferably, the process of the seventh aspect of the invention is in the pod or the anther of a plant. All preferred features of aspects one to six also apply to the seventh.

An eighth aspect of the invention provides for the use of nucleic acid according to the first to seventh aspects of the invention in the regulation/control of plant dehiscence. All preferred features of aspects one to seven also applies to the eighth.

The ninth aspect of the invention provides for the use of nucleic acid according to the first or second aspect of the invention as a probe. Such a probe can be used in techniques well known in the art to identify the presence of identical or homologous

nucleic acid sequences from any source, preferably a plant source. The ninth aspect of the invention also provides nucleic acid identified by use of the nucleic acid from aspects one or two as a probe.

5 A tenth aspect of the invention provides for the use of nucleic acid according to aspects one or two of the invention in the production of a cell, tissue, plant or part thereof, or propagating material. Again, all preferred features of aspects one and two also apply to the tenth.

10 An eleventh aspect of the invention provides for nucleic acid comprising one or more of the underlined sequences as set out in figure 1 or the primer sequences in Fig 5, Fig 9 or Fig 11. Such nucleic acid sequences are preferably used as primers in an PCR (Polymerase Chain Reaction) process in order to amplify nucleic acid sequences.

15 A twelfth aspect of the invention provides the use of nucleic acid according to the first or second aspects of the invention to identify another other protein or proteins which interact with its expression product. Such use can be carried out by the yeast two hybrid screening method (or others known in the art). The yeast two hybrid
20 screening method is described for this aspect of the invention, in general, with reference to the sequence described as DZ2. A potential way to implement the yeast 2-hybrid screen is outlined, as follows:

25 DZ2 is linked to the Gal4 DNA binding domain and expressed in yeast which contains a pGAL4-lacZ gene. For activity of lacZ a second protein is required that contains the DNA transcriptional activation domain of GAL4 and that interacts with the DZ2 protein. This is provided by making a cDNA expression library from plant DZ zones which results in fusions of plant

proteins to the GAL4 activation domain. This library is transformed into the yeast strain that contains pGAL4-LacZ and expresses the DZ2-Gal4 DNA binding domain protein fusion. Colonies that have lacZ activity are transformed with a gene for a protein that interacts with DZ2.

5

Using such a system, upstream and downstream components of any signal transduction pathway can be identified, thus resulting in further ability to control/regulate dehiscence and/or male sterility.

10 A thirteenth aspect of the invention provides for a protein, as defined according to the limitations of the second aspect of the invention (without reference to figure 1) and nucleic acid encoding the protein, wherein the protein is capable of being identified according to the use (or method) according to the twelfth aspect of the invention.

15

A fourteenth aspect of the invention provides for the use of a protein according to the fifth or sixth aspect of the invention as a probe. In this context the probe is a means to identifying interacting entities (such as other proteins), including upstream and downstream interacting signal components. A protein according to the fifth or
20 sixth aspect of the invention can be used as a probe to directly look for interactions with other proteins, i.e. purified protein can be used to look for complex formation with other plant protein, particularly isolated from the DZ zone. For example, a modified recombinant DZ2 protein can be made with a sequence tag, such as a His-tag, that enables the DZ2 + interacting protein to be directly purified on a His
25 affinity column. Alternatively, an antibody can be raised to DZ2 protein. This antibody is then used to identify DZ2 protein complexes and to purify the complexes. The DZ2 interacting proteins can be purified and microsequenced to enable cloning of the genes for these interacting proteins.

The present invention provides a particularly useful method by which plant dehiscence can be regulated/controlled.

5 In addition to the use of the present invention in the production of shatter resistance or shatter-delayed plants such as oil seed rape, the invention may be used to control/regulate pollen release (by the control/regulation of anther dehiscence) which can produce male sterile plants. The temporal and spatial expression of nucleic acid encoding a protein according to the first and second aspects of the invention may
10 require adjustment in obtaining the correct levels of dehiscence delay or prevention in different zones. For example, if pod dehiscence is required but anther dehiscence is not, it is necessary to ensure that expression of nucleic acid according to a first or second aspect of the invention has the correct temporal and spatial expression in order to obtain pod dehiscence or delay but not, to any substantial extent, anther
15 dehiscence. This can be obtained by processes known in the art and may require use of particular promoter sequences to obtain the desired result. Usually in plant transformation, some difference in the level of expression of nucleic acid is observed in different plants. In some cases, the ratio of expression levels in different tissues can vary between different plant transformants thus providing essentially tissue-
20 specific expression in one or other of the target tissues in some of the plant transformants. In the present invention, the natural expression of nucleic acid according to the first or second aspects may be predominantly higher in pod dehiscence zones and lower in the anther and funiculus dehiscence zones. However, as described above, it is possible to obtain plants in which the protein expression is
25 regulated in a particular dehiscence zone. Accordingly, a particularly useful aspect of the invention is the provision of plants which have one or both of the following features; are male sterile, are shatter resistant.

As described earlier, the process of dehiscence at the dehiscence zone involves the secretion of a number of enzymes, including hydrolytic enzymes. While previous attempts have been made to down or up regulate specific genes encoding particular proteins involved in the process of dehiscence, regulation by means of a signal transduction protein which effects expression of a number of genes is likely to be more effective than regulation of a single gene. In addition to this, the nucleic acid of the present invention has been identified as being expressed earlier than several other known genes involved in the process of plant dehiscence. This suggests that it is important earlier on in the process of plant dehiscence and can be used to control/regulate plant dehiscence at an earlier stage.

The nucleic acid encoding a signal transduction protein involved in the process of dehiscence or the signal transduction protein itself may be a component of a signal pathway that may either positively or negatively regulate pod shatter.

A more detailed explanation of such regulations/control, described with reference to a pod shatter (dehiscence) model is described below. As a skilled person will acknowledge, the model described below also relates to other general processes of dehiscence such as in the anther.

In the process of dehiscence, a particular signal transduction protein may be required to transmit a signal from the almost mature seed which initiates the expression or release of enzymes required for pod shatter. In this model, developmental signals switch on expression and/or activation of a particular signal transduction protein in the pod dehiscence zone. This leads to expression of genes required for the release of pod dehiscence zone enzymes (such as hydrolytic enzymes). In this case, prevention of activity of the signal transduction protein, for example by downregulation of expression of this protein, would result in reduced dehiscence.

Alternatively, the developing seed may transmit a signal which represses the expression and/or activity of a particular signal transduction protein until late in cell development. In this model, developmental signals switch on a particular signal
5 transduction protein which, in due course, represses the expression of genes required for release of dehiscence zone specific enzymes (such as hydrolytic enzymes). In this case, expression of a modified signal transduction protein that is constitutively active would result in reduced dehiscence.

10 A signal transduction protein which is either positively or negatively involved in the process of dehiscence can be used according to the present invention.

In addition to DZ2 several other DZ-expressed genes have been previously isolated and individually downregulated to result in *B. napus* plants that have increased
15 resistance to pod shatter; namely Sac66 (WO 96/30529 - Figure 15), DZ15 (Figure 16) and OSR 7(9) (Figure 17). It is anticipated that downregulation of more than one gene involved in pod shatter will further increase resistance to pod shatter. This could be achieved by combining different transgenes by transformation with several transgenes each designed to downregulate a different DZ-expressed gene or by
20 crossing together *B. napus* lines that individually are transformed with such transgenes. Such methods are complex either involving transformation with a construct containing multiple chimeric genes or require the maintenance of several transgenic loci in the breeding program. A preferred method is to transform with a chimeric gene consisting of a single promoter driving expression of an antisense or
25 partial sense transcript which is comprised of elements of all the DZ-expressed genes to downregulated. Similarly a single promoter could be used to drive the expression of multiple ribozymes each targeted against a different DZ-expressed gene. The use of a single promoter to drive expression of a combination of antisense, partial sense

and ribozymes is also possible. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. A suitable DZ-specific promoter would be that of DZ2, DZ2B, DZ2AT3 or ESJ2A (WO 99/13089).

5

Accordingly, the present invention provides a nucleic acid sequence according to the first or second aspects (and also all aspects which include the first or second aspects) in combination with one or more further nucleic acid sequences which are dehiscence-zone expressed. Examples of such sequences include Sac66, DZ15 and OSR(7), Figures 15-17 respectively. Such sequence may be in sense or in antisense orientation. Such a sequence may be included as full length genomic, full-length cDNA or partial sequences; the sequences may be as shown in the figures or may have the same sequence identity (both for aminoacid sequence and nucleic acid sequence) as described above for the protein according to the second aspect of the invention or the nucleic acid according to the first or second aspects of the invention. As will be recognised by those skilled in the art a partial sequence may be useful in either the sense or antisense orientation.

10

15

20

The invention is described by reference to the enclosed drawings;

Figure 1 DZ2 full length sequence showing original PCR product and primer sites

Figure 2 Amino acid alignment with bacterial response regulator proteins & *ETR1*

25

- Figure 3 Northern analysis of expression of DZ2 in pods and other tissues.
The lower panel shows the ethidium bromide-stained RNA gel prior
to blotting and probing with DZ2
- 5 Figure 4 Comparison of bacterial two-component systems with DZ2
- Figure 5 Sequence of the promoter region of *B.napus* DZ2B.
- Figure 6 Nucleic and putative peptide sequence alignments of DZ2 with DZ2B.
10
- Figure 7 Northern analysis of expression of DZ2B in pods and other tissues.
The probe was labeled DZ2B cDNA.
- Figure 8 Schematic diagram of pDZ2B-GUS-SCV
15
- Figure 9 DZ2AT3 cDNA sequence showing the putative DZ2AT3 peptide.
- Figure 10 Amino acid alignment of DZ2AT3 with DZ2 and DZ2B.
- 20 Figure 11 Sequence of the promoter region of *A.thaliana* DZ2AT3.
- Figure 12 Schematic diagram of pDZ2AT3-GUS-SCV.
- Figure 13 Schematic diagram of pPGL-DZ2as-SCV and pDZ2B-DZ2as-SCV.
25
- Figure 14 Schematic diagram of pWP357-SCV.
- Table 1 Pod shatter resistance of WP357-SCV transformants.

Figure 15 Nucleic acid sequence and putative amino acid sequence of Sac66.

Figure 16 Nucleic acid sequence and putative amino acid sequence of DZ15.

Figure 17 Nucleic acid sequence and putative amino acid sequence of OSR7 (9)

The present invention is now described with reference to the following, non-limiting examples.

Example 1- Isolation and characterisation of expression of DZ2

Plant Material

Seeds of *B. napus* cv Rafal were grown as described by Meakin and Roberts, (1990a) with the following modifications. Single seedlings were potted into 10cm pots, and after vernalization, were re-potted into 21cm pots. At anthesis tags were applied daily to record flower opening. This procedure facilitated accurate age determination of each pod. Pods were harvested at various days after anthesis (DAA). The dehiscence zone was excised from the non-zone material and seed using a scalpel blade (Meakin and Roberts (1990b)) and immediately frozen in liquid N₂ and stored at -70°C.

RNA Isolation

All chemicals were molecular biology grade and bought from either Sigma Chemical Ltd (Dorset, UK), or Fisons (Loughborough, UK). Total RNA was extracted using the polysomal extraction method of Christoffersen and Laties, (1982), with the following alterations. The plant material was ground to a powder in liquid N₂ and then in 10 volumes of extraction buffer (200mM Tris-acetate [pH 8.2], 200mM magnesium

acetate, 20mM potassium acetate, 20mM EDTA, 5% w/v sucrose, after sterilisation 2-mercaptoethanol was added to 15mM and cycloheximide added to a final concentration of 0.1 mg ml⁻¹). The supernatant was then layered over 8 ml 1M sucrose made with extraction buffer and centrifuged in a KONTRON™ (Switzerland) TFT 70.38 rotor at
5 45,000rpm (150,000g) for 2 hr at 2°C in a Kontron CENTRIKON™ T-1065 ultra-centrifuge. Pellets were then resuspended in 500µl 0.1M sodium acetate, 0.1% SDS, pH 6.0 and phenol/chloroform (1:1 v/v) extracted and the total RNA precipitated. Poly(A)⁺ RNA was isolated from total RNA extracted, from both the zone and non-zone tissue of 40, 45 and 50 DAA pods, using a Poly(A) QUIK™ mRNA purification kit
10 (Stratagene, Cambridge, UK) following the manufacturers instructions, and then bulked together. Total RNA was also extracted from leaves, stems, seeds and pods using a method described by Dean *et al.*, (1985) for use in Northern analyses.

Differential display

15

This was performed essentially as described by Liang and Pardee (1992) using RNA extracted from 40 DAA pod dehiscence zones and non-zones. First strand cDNA copies of the RNAs (40 DAA DZ/NZ) were made using 50U M-MLV (Moloney Murine Leukemia Virus) reverse transcriptase (50U/µL) (Stratagene) in a 20µL
20 reaction containing 1x M-MLV buffer, 2.5mM dNTPs (Pharmacia), 1µg RNA, 30U RNase inhibitor (Promega) and 10µM oligo dT anchor primer 7 (5'-TTTTTTTTTTTTTTGG-3'). The reaction conditions were as follows: 65°C for 5 minutes, 37°C for 90 minutes and 95°C for 5 minutes. Following first strand cDNA synthesis, 60µL dH₂O were added and the samples were either used directly for
25 PCR or stored at -20°C.

For PCR, 2µL cDNA were used as template in a 20µL reaction containing 1x PCR buffer, 1mM MgCl₂, 2µM dNTPs, 10µM oligo dT anchor primer 7 (5'-

TTTTTTTTTTTTTGG-3'), 2.5 μ M arbitrary primer A (5'- AGC CAG CGA A -3'), 0.5 μ L 35S-dATP (> 1000 Ci/mmol) (Amersham) and 1U *Taq* DNA polymerase (5U/ μ L) (Gibco BRL). The thermocycling conditions were as follows: 40 cycles of 94°C for 30 seconds, 40°C for 2 minutes, 72°C for 30 seconds followed by 72°C for 5 minutes. The PCR products were fractionated on a 5 % polyacrylamide/7M urea gel after addition of 5 μ L loading buffer (95 % (v/v) formamide, 20mM EDTA, 0.05 % (w/v) xylene cyanol, 0.05 % (w/v) bromophenol blue) to each sample. Following electrophoresis the gel was dried at 80°C under vacuum for 1 hour then exposed to X-ray film (BioMax-MR, Kodak) in a light tight cassette for 48 hours.

The dried gel and autoradiogram were aligned so that bands that appeared in the DZ and not in NZ could be cut out and the DNA eluted according to Liang et al. (1995). The eluted PCR products (4 μ L) were reamplified in a 40 μ L reaction containing 1x PCR buffer, 1mM MgCl₂, 20 μ M dNTPs, 10 μ M oligo dT anchor primer 7 (5'- TTTTTTTTTTTTTTGG-3'), 2.5 μ M arbitrary primer A (5'- AGC CAG CGA A -3') and 2U *Taq* DNA polymerase (5U/ μ L) (Gibco BRL) using the following thermocycling conditions: 40 cycles of 94°C for 30 seconds, 40°C for 2 minutes, 72°C for 30 seconds followed by 72°C for 5 minutes. The resulting PCR product was cloned into the TA cloning vector (Invitrogen) and sequenced (Figure 1). In order to prepare an antisense strand-specific riboprobe, the PCR product was subcloned into pBluescript (Stratagene).

Expression analysis and characterisation of DZ2

Northern analysis using an antisense strand-specific riboprobe to the DZ2 PCR product, showed that DZ2 hybridised to a transcript of 0.6kb which is expressed in the DZ of 20-50 DAA pods with a peak in expression at 40DAA. Minimal expression was observed in the pod NZ [Figure 2]. A random-primed labelled DNA probe (Stratagene) of the 330bp DZ2 PCR product (amplified using primers DZ2FL

and DZ2RL - see Figure 1) was used to screen a *B. napus* DZ cDNA library from which, following three rounds of screening to obtain pure plaques, a full length DZ2 cDNA (606bp) was obtained (Figure 1). An antisense strand-specific riboprobe of the full length DZ2 cDNA was hybridised to total RNA extracted from pod DZ/NZ (as in Figure 2), leaf abscission zones (AZ) and non-zones (NZ) (following exposure to 10 μ L/L ethylene for 72 hours), seed, root, flower and leaf. Figure 3 shows that DZ2 hybridises to a 0.6kb message which is present in the pod DZ at 20-50 DAA with maximum expression at 40DAA. Again there is minimal expression in pod NZ and no apparent expression of DZ2 in AZ, NZ, leaf, root, seed or flower RNA. By the sensitive technique of RT-PCR analysis DZ2 expression can also be detected in anthers and the funiculus, both tissues that contain dehiscent zones

The 606bp cDNA (DZ2) encodes a putative protein of 136 amino acids. Comparison of the DZ2 translated sequence to the OWL protein database [Bleasby and Attwood (1994)] showed low but consistent homology to a group of bacterial proteins comprising two-component regulatory systems. In particular, DZ2 possesses the conserved amino acid residues required for phosphorylation of the receiver domain of the response regulator component (see Figure 4). DZ2 plays a role in a signal transduction cascade resulting at least in one respect in pod shatter. It is therefore a good candidate for down-regulation of pod shatter processes using antisense technology. DZ2 is a novel plant protein in that independent proteins with homology to bacterial receivers are yet to be reported in plants.

The full length cDNA was excised from the pBluescript cloning vector by digestion with *Eco*RI and *Xho*I restriction enzymes (Gibco BRL). Following purification from a 1% agarose gel the 606bp cDNA was random primed labelled (Stratagene) and used to screen a *B. napus* genomic library in the BlueStar vector. Following three rounds of screening to obtain pure plaques, a single genomic clone was isolated

which carries a 15kb genomic DNA insert. The promoter of the DZ2 gene is isolated from this genomic clone using standard techniques (see Example 2).

Example 2 - Isolation and characterisation of the *B.napus* DZ2B promoter.

To obtain the *B.napus* DZ2 promoter a *B.napus* genomic library was screened with a labelled DZ2 probe. The full length cDNA was excised from the pBluescript cloning vector (Stratagene) by digestion with *Eco*RI and *Xho*I restriction enzymes (Gibco BRL). Following purification from a 1% agarose gel the 606bp cDNA was random primed labelled (Stratagene) and used to screen a *B. napus* genomic library in the BlueStar vector (Novagen). Following three rounds of screening to obtain pure plaques, a single genomic clone was isolated which carries a 15kb genomic DNA insert. The region hybridising to DZ2 was sequenced and found to encode a protein homologous to, but not identical to DZ2. This DZ2-like gene was designated DZ2B (Figure 5). The primers DZ2BFL (Figure 5) and T7 were used to PCR out a DZ2B cDNA from the *B.napus* DZ cDNA library.

5' AACCAAGTCAGTAGAAGTG 3' DZ2BFL

5' AATACGACTCACTATAGG 3' T7

The DZ2 and DZ2B cDNAs are 80% identical (according to the default parameters of the GAP computer program, version 6, Deveraux *et al.*, 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) at the nucleotide level in the region of overlap of the coding sequences (Figure 6a) and the putative proteins encoded by DZ2 and DZ2B are 80% identical (according to the default parameters of the GAP computer program, version 6, Deveraux *et al.*, 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) (Figure 6b). Sequence analysis of more DZ2 and DZ2-like cDNAs and Southern analysis shows that there are two DZ2 genes in *B.napus*, DZ2 and DZ2B,

each represented by 2 slightly different cDNAs. This is consistent with one gene being encoded by the *B.campestris* derived-genome and the other from the genome derived from *B.oleracea*.

5 RT-PCR with primers specific to DZ2B showed that DZ2B is only expressed in pods. This was confirmed by northern analysis which showed preferential expression in the DZ (Figure 7). Thus DZ2B has a similar pattern of expression as DZ2 and is thus a suitable source of a DZ-expressed promoter.

10 Primers DZ2BGenF and DZ2BGenR were used to PCR a 1253bp DZ2B promoter fragment (Figure 5).

5' GGCTCTAGACGAACTGCGGAGCAAGG 3' DZ2BGENF

5' CTGCCATGGTCGGTTTTTTTTTCTTCGAAC 3' DZ2BGENR

15

These primers introduced an *Xba*I site at the 5' end of the PCR fragment and an *Nco*I site around the initiating Met of DZ2B. Thus the PCR fragment was cloned as an *Xba*I, *Nco*I fragment between the *Xba*I and *Nco*I sites of pWP272 (WO 99/10389) forming pDZ2B-GUS. The chimeric pDZ2B-GUS-CaMV polyA gene was then transferred as an *Xba*I, *Xho*I fragment between the *Xba*I and *Sal*I sites of pSCV nos-nptII (WO 95/20668) forming pDZ2B-GUS-SCV (Figure 8). The pDZ2B-GUS-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed *B.napus* plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989). Gus expression is observed in the pod DZ.

25

Example 3 - Isolation and characterisation of a DZ2 *Arabidopsis thaliana* homologue

To demonstrate that a DZ2 orthologous gene can be isolated from another plant species the functional equivalent of B.napus DZ2 / DZ2B was isolated from
5 Arabidopsis thaliana. The B.napus DZ2 cDNA was used as a probe to screen an Arabidopsis cDNA library (J. Giraudat, ISV-CNRS, France). Figure 9 shows the sequence of a cDNA (DZ2AT3) that hybridised to the DZ2 probe. DZ2AT3 has 85% nucleic acid identity to DZ2 and 85% to DZ2B (according to the default
10 parameters of the GAP computer program, version 6, Deveraux *et al.*, 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) in the coding regions which are common to all three sequences. The putative peptide encoded by DZ2AT3 has 80% identity to DZ2 and 80% to DZ2B (according to the default parameters of the GAP computer program, version 6, Deveraux *et al.*, 1984,
15 and available from the University of Winsconsin Genetics Computer Group (UWGCG)) in the regions which are common to all three sequences (Figure 10). RT-PCR analysis of RNA isolated from leaves, roots, flowers and siliques showed that DZ2AT3 was specifically expressed in siliques. Southern hybridisation analysis showed that the DZ2AT3, DZ2 and DZ2B probes each identify a single identical
20 band in A.thaliana. This indicates that A.thaliana contains one DZ2 gene in contrast to B.napus which contains two.

The Genome walker kit (Clontech) was used to isolate the DZ2AT3 promoter from A.thaliana genomic DNA. Nested PCR was performed using primer GW1 first, then
25 AT3GW2 each in conjunction with the Genome Walker kit primer (Figure 9). Figure 11 shows the sequence of the promoter region of DZ2AT3 thus obtained. The primers ATDZ2F and ATDZ2R were used to PCR a 1195bp promoter fragment from the DZ2AT3 genomic sequence (Figure 11).

5' CACTAGTAGGGCACGCGTGGTCG 3' ATDZ2F
 5' TCCATGGTCGATTTCTTTTCTCTCAAG 3' ATDZ2R

These primers introduced an SpeI site at the 5' end of the PCR fragment and an
 5 NcoI site around the initiating Met of DZ2AT3. Thus the PCR fragment was cloned
 as an SpeI, NcoI fragment between the XbaI and NcoI sites of pWP272 (WO
 99/13089) forming pDZ2AT3-GUS. The chimeric pDZ2AT3-GUS-CaMV polyA
 gene was then transferred as a SalI, XhoI fragment into the SalI site of pSCV nos-
 nptII (WO 95/20668) forming pDZ2AT3-GUS-SCV (Figure 12). The pDZ2AT3-
 10 GUS-SCV binary vector was transferred to the agrobacterial strain pGV2260 and
 transformed *B.napus* plants produced by agrobacterial transformation essentially as
 described in Moloney M et al., (1989). Gus expression is observed in the pod DZ.

**Example 4 - Production of shatter-resistant *B.napus* plants by antisense
 15 downregulation of DZ2**

Downregulation of the DZ2 gene or reduction in DZ2 protein levels in the pod DZ
 will result in plants that are resistant (or more resistant than without this
 modification) to pod shatter. Standard techniques, commonplace in the art, such as
 20 the expression of antisense DZ2 mRNA, full sense mRNA, partial sense mRNA or a
 ribozyme directed against DZ2 mRNA are effective. Expression of these RNAs
 requires a promoter that is active in the pod DZ at the time at which DZ2 is
 expressed. Ideally the promoter will be pod DZ-specific, however a useful promoter
 may be pod-specific or even constitutively active. A suitable promoter would be that
 25 of DZ2. Although DZ2 is expressed in the anther DZ, pod DZ and funiculus DZ,
 DZ2 promoter -GUS fusion studies show that in different transformants the relative
 level of expression in these three sites is variable but is stability heritable. Thus
 some transformants are obtained in which expression is largely or exclusively

confined to the pod DZ. This suggests that the pDZ2 promoter is comprised of distinct elements each specifying expression in a particular DZ. Alternatively the site of transgene integration may influence relative expression levels in the DZ tissues. The DZ2 promoter is therefore linked to the DZ2 cDNA such that the DZ2 is in the antisense orientation forming pDZ2-antiDZ2. This chimeric gene is transferred to the binary vector pNos-NptII-SCV (W0 96/30529). This binary vector is transferred to the agrobacterial strain pGV2260 and transformed *B.napus* plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989) Plant Cell Reports 8, 238-242. A proportion of transformed *B.napus* plants exhibit reduced levels of DZ2 message and are resistant to pod shatter.

Example 5 - Production of shatter-resistant *B.napus* plants by antisense downregulation of DZ2

Downregulation of the DZ2 gene or reduction in DZ2 protein levels in the pod DZ will result in plants that are resistant to pod shatter. Techniques such as the expression of antisense DZ2 mRNA, full sense mRNA, partial sense mRNA or a ribozyme directed against DZ2 mRNA will be effective. Expression of these RNAs requires a promoter that is active in the pod DZ at the time at which DZ2 is expressed. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. Although DZ2 / DZ2B is expressed in the anther DZ, pod DZ and funiculus DZ, DZ2B promoter -GUS fusion studies show that in different transformants the relative level of expression in these three sites is variable but is stably heritable. Thus some transformants are obtained in which expression is largely or exclusively confined to the pod DZ. This suggests that the pDZ2 promoter is comprised of distinct elements each specifying expression in a particular DZ. Alternatively the site of transgene integration may influence relative expression levels in the DZ tissues. Thus a suitable DZ-specific promoter would be that of DZ2, DZ2B, DZ2AT3 or ESJ2A (WO 99/13089).

The primers DZ2FLA and DZ2RLA were used to PCR a 349bp fragment from the DZ2 cDNA:-

5 5' GGCGAATTCCGGTGAGGAGGCAGTAATC 3' DZ2FLA
 5' GGCCCATGGCATAACATACACACTTAGAC 3' DZ2RLA

10 The primers introduce an EcoRI and NcoI site at the ends of the DZ2 PCR fragment. To link the DZ2 PCR fragment in an antisense orientation to the promoter of ESJ2A (pPGL) the DZ2 PCR fragment was cloned as a NcoI, EcoRI fragment between the NcoI and EcoRI sites of pWP272 (WO 99/13089) forming pPGL-DZ2as. The pPGL-antisense DZ2 chimeric gene was transferred as a XbaI, XhoI fragment from pDZ2as into the XbaI and SalI sites of the binary vector pSCV nos-nptII (WO 95/20668) forming pPGL-DZ2as-SCV (Figure 13a).

15 The pPGL-DZ2as-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed B.napus plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989). A proportion of transformed B.napus plants exhibit reduced levels of DZ2 and DZ2B message and were resistant
 20 to pod shatter

25 Similarly, to link the DZ2 PCR fragment in an antisense orientation to the promoter of DZ2B, the DZ2 PCR fragment is cloned as a NcoI, EcoRI fragment between the NcoI and EcoRI sites of pDZ2B-GUS forming pDZ2B-DZ2as. The pDZ2B-DZ2as chimeric gene is transferred as a XbaI, XhoI fragment from pDZ2B-DZ2as into the XbaI and SalI sites of the binary vector pSCV nos-nptII (WO 95/20668) forming pDZ2B-DZ2as-SCV (Figure 13b).

The pDZ2B-DZ2as-SCV binary vector is transferred to the agrobacterial strain pGV2260 and transformed *B.napus* plants. Again a proportion of transformed *B.napus* plants exhibit reduced levels of DZ2 and DZ2B message and are resistant to pod shatter.

5

Similarly a proportion of *B.napus* plants transformed with a pDZ23A-DZ2as-SCV construct exhibit reduced levels of DZ2 and DZ2B message and are resistant to pod shatter.

10 **Example 6 - Production of shatter-resistant *B.napus* plants by antisense downregulation of multiple DZ-expressed genes**

In addition to DZ2 several other DZ-expressed genes have been previously isolated and individually downregulated to result in *B.napus* plants that have increased resistance to pod shatter; namely Sac66 (WO 96/30529 Figure 15), DZ15 (Figure 16) and OSR 7(9) (Figure 17). It is anticipated that downregulation of more than one gene involved in pod shatter will further increase resistance to pod shatter. This could be achieved by combining different transgenes by transformation with several transgenes each designed to downregulate a different DZ-expressed gene or by crossing together *B.napus* lines that individually are transformed with such transgenes. Such methods are complex either involving transformation with a construct containing multiple chimeric genes or require the maintenance of several transgenic loci in the breeding program. A preferred method is to transform with a chimeric gene consisting of a single promoter driving expression of an antisense or partial sense transcript which is comprised of elements of all the DZ-expressed genes to be downregulated. Similarly a single promoter could be used to drive the expression of multiple ribozymes each targeted against a different DZ-expressed gene. The use of a single promoter to drive expression of a combination of

antisense, partial sense and ribozymes is also possible. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. A suitable DZ-specific promoter would be that of DZ2, DZ2B, DZ2AT3 or ESJ2A.

5

Consequently the ESJ2A promoter was linked to a multiple antisense gene consisting of elements of Sac66, DZ2, DZ15 and OSR 7(9) in the following manner:- The original DZ15 PCR product in pCRII (Invitrogen) (see Figure 16) was cloned as an EcoRI fragment into pBluescript SK (Stratagene) forming pDZ15-BS, such that the DZ15 3' end is nearest the SstI site of the vector. T7 and DZ15RL primers were used to PCR a 456bp DZ15 fragment from pDZ15-BS which was cloned into the EcoRV site of pGEM5zf (Promega) forming pWP351, such that the DZ15 3' end is nearest the SphI site of the vector.

10

15

5' AATACGACTCACTATAGG 3' T7

5' AACAGCTGAAAACCTCACGAAG 3' DZ15RL

The EcoRI, NcoI fragment of pWP351 cloned between the EcoRI and NcoI sites of pDZ2-BS forming pWP356. pDZ2-BS consists of the DZ2 cDNA cloned as an EcoRI, XhoI fragment into pBluescript SK such that the 3' end is nearest the KpnI site of the vector. A 361bp Sac66 fragment was PCR'd from the Sac66 cDNA (WO 96/30529) using the primers F1 and RI which introduce NcoI and PstI sites into the ends of the PCR product.

20

25

5' GGCCCATGGCTGCCAAGCTTTGAGTAGC 3' F1

5' GGCCTGCAGTGCCTAGGATCTGGAAGC 3' RI

The Sac66 PCR product was cloned as an NcoI, EcoRI fragment between the NcoI and EcoRI sites of pWP272 (WO 99/13089) forming pWP288A. EcoRI DZ15 + DZ2 and OSR 7(9) fragments from pWP356 and pOSR 7(9)-CRII were cloned into the EcoRI site of pWP288A such that DZ15 + DZ2 and OSR 7(9) are in an antisense orientation with respect to PGL promoter. (pOSR 7(9)-CRII consists of the 306bp OSR 7(9) PCR fragment (see Figure 17) cloned into pCRII (Invitrogen)). The chimeric pPGL-antisense Sac66 + DZ2 + DZ15 + OSR 7(9) gene was transferred as a XbaI, XhoI fragment into the XbaI and SalI sites of the binary vector pSCV nos-nptII (WO 95/20668) forming pWP357-SCV (Figure 14). The pWP357-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed *B.napus* plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989) Plant Cell Reports 8, 238-242.

Resistance to podshatter was measured using an impact pendulum device (Liu X-Y, Macmillan RH and Burrow RP 1994 Journal of Texture Studies 25 p179-189) (Table1). The mean energy values shown in Table 1 represent the energy required to rupture the pod on impact with the pendulum. These values are an average from measurements of 20 mature pods. The letters A to L indicate grouping of transformants with significantly different podshatter resistance (ie Group A is significantly different from B when analysed by ANOVA using a Fisher PLSD analysis with a significance level of 95% (Statview 512+)). Lines with a number of letters are not significantly different from other lines sharing the same letter. The results shown in Table 1 indicate that 24 lines exhibited significantly higher resistance to podshatter than non-transformed controls whilst 17 lines were not significantly different from the control. The degree and frequency of pod shatter resistance achieved with pWP357-SCV was greater than that obtained by transformation with constructs that downregulate a single DZ-expressed gene.

Table 1

| PLANT ID | | | | | |
|-------------|-------|---|---|---|---|
| MEAN ENERGY | | | | | |
| A213-24 | 6.752 | A | | | |
| A213-53 | 5.203 | B | | | |
| A213-34 | 3.864 | C | | | |
| A213-21 | 3.673 | C | | | |
| A213-4 | 3.54 | C | | | |
| A213-61 | 3.516 | C | | | |
| A214-30 | 3.46 | C | | | |
| A214-27 | 3.397 | C | | | |
| A213-8 | 3.277 | C | | | |
| A213-70 | 3.271 | C | D | | |
| A214-13 | 3.182 | C | D D E | | |
| A213-11 | 3.182 | C | D D E F | | |
| A213-69 | 3.005 | C | D D E F G | H | |
| A213-60 | 2.945 | | D D E F G H | I | J |
| A214-7 | 2.843 | | D D E F G H I | J | K |
| A213-64 | 2.687 | | D D E F G H I J | K | L |
| A214-14 | 2.613 | | D D E F G H I J K | L | M |
| A213-28 | 2.581 | | D D E F G H I J K L | M | N |
| A213-9 | 2.547 | | D E F G H I J K L M N | O | P |
| A213-42 | 2.442 | | E F G H I J K L M N O P Q R S T U V W X Y Z | | |
| A213-31 | 2.431 | | F G H I J K L M N O P Q R S T U V W X Y Z | | |
| A214-10 | 2.42 | | G H I J K L M N O P Q R S T U V W X Y Z | | |
| A213-38 | 2.295 | | H I J K L M N O P Q R S T U V W X Y Z | | |
| A214-8 | 2.26 | | I J K L M N O P Q R S T U V W X Y Z | | |
| A213-19 | 2.213 | | J K L M N O P Q R S T U V W X Y Z | | |
| A214-25 | 2.128 | | K L M N O P Q R S T U V W X Y Z | | |
| A213-38 | 2.059 | | L M N O P Q R S T U V W X Y Z | | |
| A213-16 | 2.005 | | M N O P Q R S T U V W X Y Z | | |
| A213-63 | 1.928 | | N O P Q R S T U V W X Y Z | | |
| A213-33 | 1.91 | | O P Q R S T U V W X Y Z | | |
| A213-32 | 1.901 | | P Q R S T U V W X Y Z | | |
| A213-37 | 1.791 | | Q R S T U V W X Y Z | | |
| RV27CONT | 1.787 | | R S T U V W X Y Z | | |
| A213-10 | 1.623 | | S T U V W X Y Z | | |
| A213-58 | 1.595 | | T U V W X Y Z | | |
| A214-24 | 1.55 | | U V W X Y Z | | |
| A213-3 | 1.518 | | V W X Y Z | | |
| A213-27 | 1.495 | | W X Y Z | | |
| A213-40 | 1.395 | | X Y Z | | |
| A213-47 | 1.315 | | Y Z | | |
| A213-43 | 1.286 | | Z | | |
| A213-30 | 1.241 | | | | |

REFERENCES

- 5 1. Bleasby A J and Attwood T K, OWL – A Non-redundant Composite Protein
Sequence Database. *Nucleic Acid Research* 22:3574-3577 (1994)
2. Christoffersen and Laties, *Proc. Natl. Acad. Sci.* 79, 4060-4063 (1982)
- 10 3. Dean *et al.*, *EMBO J.* 4: 3055-3061 (1985)
4. Deveraux *et al.*, *Nucl. Acids Res.* 12:387 (1984)
- 15 5. Hocking P J and Mason L: Accumulation, distribution and redistribution of
dry matter and mineral nutrients in fruits of canola (oilseed rape) and the
effect of nitrogen fertiliser and windrowing. *Australian Journal of*
Agriculture Research 44: 1377-1388 (1993)
- 20 6. Hakwa *et al.*, *Science* 198:105-63 (1977)
7. Jenkins *et al.*: 5th International Conference of Plant Molecular Biology,
Abstract 310 (1997)
- 25 8. Johnson-Flanagan AM and Spencer MS: Ethylene production during
development of mustard (*Brassica juncea*) and canola (*Brassica napus*) seed.
Plant Physiol 106: 601-606 (1994)
9. Liang P and Pardee A: Differential display of eukaryotic messenger RNA by

means of the polymerase chain reaction. Science 257: 967-971 (1992)

- 5
10. Laing P, Bauer D, Averboukh L, Warthoe P, Rohrwild M, Muller H, Strauss M, Pardee A B: Analysis of altered gene expression by differential display. Methods in Enzymol 254: 304-321 (1995)
11. Meakin P J and Roberts J A: Dehiscence of fruit in oilseed rape (*Brassica napus* L.): anatomy of pod dehiscence. J Expt. Bot 41: 995-1002 (1990a)
- 10 12. Meakin P J and Roberts J A: Dehiscence of fruit in oilseed rape (*Brassica napus* L.): the role of cell wall degrading enzymes and ethylene. J Expt. Bot 41: 1003-1011 (1990b)
13. Moloney M *et al*; Plant Cell Reports, 8, 238-242 (1989)
- 15
14. Needleman and Wunsch J. Mol. Biol. 48:443 (1970)
15. Smith and Waterman Adv. Appl. Math 2: 482 (1981)

CLAIMS

1. Nucleic acid encoding a signal transduction protein involved in the process of dehiscence.
- 5
2. Nucleic acid as claimed in claim 1 wherein the process involves the production of a hydrolytic enzyme.
3. Nucleic acid as claimed in claim 1 or claim 2 which is naturally expressed in a dehiscence zone.
- 10
4. Nucleic acid encoding a protein wherein the protein:
- a) comprises the amino acid sequence shown in Figure 1 or;
- 15
- b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above and has at least 40% amino acid sequence identity therewith; or
- 20
- c) is a fragment of a protein as defined in a) or b) above, which is at least 10 amino acids long.
5. Nucleic acid as claimed in any one of claims 1 to 4 which comprises the sequence set out in Figure 1 or a fragment thereof which is at least 30 bases long.
- 25
6. Nucleic acid, as claimed in any one of claims 1 to 5 in combination with one or more further nucleic acid sequence which is dehiscence-zone expressed.

7. Nucleic acid which is antisense to nucleic acid as claimed in any one of claims 1 to 6.
- 5 8. Nucleic acid as claimed in any one of claims 1 to 7 including a promoter or other regulatory sequence which controls expression of the nucleic acid.
9. Nucleic acid which is the naturally occurring promoter or other regulatory sequence which controls expression of nucleic acid as claimed in any one of
10 claims 1 to 8.
10. Nucleic acid as claimed in any one of claims 1 to 9 which is in the form of a vector.
- 15 11. A cell comprising nucleic acid as claimed in any one of claims 1 to 10.
12. A plant cell as claimed in claim 11.
13. A process for obtaining a cell as claimed in claim 11 or claim 12 comprising
20 introducing nucleic acid as claimed in any one of claims 1 to 10 into said cell.
14. A plant or a part thereof comprising a cell as claimed in claim 11 or claim
25 12.
15. Propagating material or a seed comprising a cell as claimed in claim 11 or claim 12.

16. A process for obtaining a plant or plant part as claimed in claim 14 or claim 15 comprising obtaining a cell as claimed in claim 11 and growth thereof or obtaining a plant, plant part, or propagating material as claimed in claim 14 or claim 15 and growth thereof.
- 5
17. A signal transduction protein involved in the process of plant dehiscence.
18. A protein which:
- 10
- a) comprises the amino acid sequence shown in Figure 1 or;
- b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above, and has at least 40% amino acid sequence identity therewith; or
- 15
- c) a fragment of a protein as defined in a) or b) above which is at least 10 amino acids long.
19. A protein as claimed in claim 17 or claim 18 which is isolated or
- 20
- recombinant.
20. A process for regulating/controlling dehiscence in a plant or a part thereof, the process comprising obtaining a plant or part thereof as claimed in claim 14.
- 25
21. A process as claimed in claim 20 which comprises obtaining a plant cell as claimed in claim 21 or part of a plant as claimed in claim 14 and deriving a plant therefrom.

22. A process as claimed in claim 20 which comprises obtaining propagating material or a seed as claimed in claim 15 and deriving a plant therefrom
- 5 23. A process as claimed in claim 20 wherein the dehiscence is of a pod or of an anther.
24. Use of nucleic acid as claimed in any one of claims 1 to 10 in the regulation/control of plant dehiscence.
- 10 25. Use of nucleic acid as claimed in any one of claims 1 to 10 as a probe.
26. Use of nucleic acid as claimed in any one of claims 1 to 10 in the production of a cell, tissue, plant part thereof or propagating material.
- 15 27. Nucleic acid comprising one or more of the underlined sequences as set out in Figure 1, or one or more of the primer sequences in Figure 5, 9 and/or 11.
28. Use of the nucleic acid as claimed in claim 27 as a PCR primer.
- 20 29. Use of a protein as claimed in any one of claims 17 to 19 as a probe.

FIG. 1

NUCLEIC ACID AND PREDICTED PROTEIN SEQUENCE OF DZ2

1/27

NcoI

```

1  GGCACGAGCAGAATCGAAGATGGCAACAAATCCATGGGAGATATCGAGAAATAAGAA 60
   M A T K S M G D I E K I K K

61  GAACTAAACGTGTGATCGTCGATGATCCACTAAACCTTATAATCATGAGAAGAT 120
   K L N V L I V D D D P L N L I I H E K I

121 CATCAAAGCGATTGGGGTATTTACAGACAGCGGAATAACGGTGAGGAGGCAGTAATCAT 180
   I K A I G G I S Q T A N N G E E A V I I
       ↓
       DZ2FL→

181 CCACCGTGACGGCGGCTCATCTTTTGACCTTATCCTAATGGATAAAGAAATGCCCGAGAG 240
   H R D G G S S F D L I L M D K E M P E R

241 GGATGGTGTTCGACAACCTAAGAAGCTAAGAGAAATGGAAGTGAAGTCAATGATTGTTGG 300
   D G V S T T K K L R E M E V K S M I V G

301 GGTGACTTCACTGGCTGACAATGAAGAGGAGCGCAGGCTTTCATGGAAGCTGGACTTAA 360
   V T S L A D N E E E R R A F M E A G L N
  
```

2 / 27

```

361 CCATTGCTTGGCAAAACCGTTAACCAAGGACAAGATCATCCCTCTCATTAACCAACTCAT 420
    H C L A K P L T K K D K I I P L I N Q L M
421 GGATGCTTGATGGATATATATTTATATATGGAACACACACATAATAACGCTCTAAGTGTG 480
    D A *
481 TATGTATGCATAGATACTTGCATGTGTGTGTTTAGAATTAGGGTTCCTTATCGTCCGT 540
    HindIII
541 GATATATAATCATGTAAGTTGTTGCTTTAAGCTTATAAAATATTTAAATAAGGGTTTCCT 600
601 CTACC

```

The primer sites for DZ2FL and DZ2RL are underlined, as are the recognition sequences for *Nco*I and *Hind*III restriction enzymes. Shown in bold are the conserved amino acid residues required for phosphorylation. The extent of the original PCR product isolated by differential display is shown by ↓.

FIG. 1 CONT'D

3/27

Alignment of the predicted protein sequence of DZ2 with those of
bacterial response regulator proteins.

| | | | | | |
|-------|-------------------|--|-------------------------------------|-------------------|-------------------|
| | 1 | | | | 50 |
| DZ2 | MATKSMGDIE | KIKKKLNVLI | VDDDPLNLII | HEKIIKAIG | GISQTANNGE |
| OMPR | | .MQENYKILV | VDDDMRLRAL | LERYLTEQGF | .QVRSVANAE |
| PHOB | | ...MARRILV | VEDEAPIREM | VCFVLEQNGF | .QPVEAEDYD |
| NTRC | | ..MQRGIVWV | VDDDSSIRWV | LERALAGAGL | .TCTTFENG |
| SPOOF | | ..MMNEKILI | VDDQYGIRIL | LNEVFNKEGY | .QTFOAANGL |
| CHEY | | MADKELKFLV | VDDFSTMRI | VRNLLKELGF | NNVEEAEDGV |
| ETR | |LKVLV | MDENGVSVMV | TKGLLVHLGC | EVTTVSSNEE |
| | 51 | | | | 100 |
| DZ2 | EAVIIHRDGG | SSFDLILMDK | EMPERDGVST | TKKLREMEVK | SM..IVGVTS |
| OMPR | QMDRLLTR.. | ESFHLMVLDL | MLPGEDGLSI | CRRLRSQS.. | NMPPIIMVTA |
| PHOB | SAVNQLNE.. | PWPDILLLDW | MLPGGSGIQF | IKHLKRESMT | RDIPVVMLTA |
| NTRC | EVLAALAS.. | KTPDVLLSDI | RMPGMDGLAL | LKQIKQ..RH | PMLPVIIMTA |
| SPOOF | QALDIVTK.. | ERPDVLVLLD | KIPGMDGIEI | LKRMKV..ID | ENIRVIIMTA |
| CHEY | DALNKLQA.. | GGYGFVISDW | NMPNMDGLEL | LKTIRADGAM | SALPVLMTA |
| ETR |CLRVS | HEHKVVFMDV | CMPGVENYQI | ALRI..... | .HXPLLVALS |
| | 101 | | | | 150 |
| DZ2 | LADNEEERRA | FMEAGLNHCL | AKPLTKDKII | PLINQLMDA | |
| OMPR | KGEEVDRIVG | .LEIGADDYI | PKPFNPPELL | ARIRAVLRRQ | ANELPGAPS. |
| PHOB | RGEEEDRVRG | .LETGADDYI | TKPFSPKELV | ARIKAVMRRI | SPM..... |
| NTRC | HSDLDAVSA | .YQQGAFDYL | PKPFDIDEAV | ALVERAISHY | QEQQQPRNIE |
| SPOOF | YGELDMIQES | .KELGALTHF | AKPFDIDEIR | DAVKKYLPLK | SN..... |
| CHEY | EAKKENIIAA | .AQAGASGYV | VKPFTPATLE | EKLNKIFEKL | GM..... |
| ETR | GNTDKSTKEK | CMSFGLDGV | LKPVSLDNIR | DVLSDLL... | |
| | 151 | | | | |
| DZ2 | | | | | |
| OMPR | | OmpR = <i>E. coli</i> | (Involved in osmoregulation) | | |
| PHOB | | PhoB = <i>E. coli</i> | (Involved in phosphate utilisation) | | |
| NTRC | VNGP | NtrC = <i>S. typhimurium</i> | (Involved in nitrogen utilisation) | | |
| SPOOF | | SpoOF = <i>B. subtilis</i> | (Involved in sporulation) | | |
| CHEY | | CheY = <i>E. coli</i> | (Involved in chemotaxis) | | |
| ETR | | ETR = <i>A. thaliana ETR1</i> gene encoding an ethylene receptor | (partial amino acid sequence) | | |

The predicted protein sequence of DZ2 is shown in bold as are the conserved amino acid residues required for phosphorylation of the protein

FIG. 2

4 / 27

Expression analysis of DZ2 in various plant organs using Northern




FIG. 3

5 / 27

Comparison of bacterial two-component regulatory systems with DZ2




FIG. 4

FIG. 5

TATATAAATACGGTTTAAACAGATATGTTCTGTTATAAATGTAATTCNATGTGCCNNTCAANITTTATTTTNAATTNGT 78
 TNTACTAGGACATTAGTTTTAACNITTTTATATATCATGTACAAAAAANNTTTTATATNTCAACTATGA 156
 GCAATTATCTTATAGTGTTCCTTTCCAGAAATTTGACGACAACCTAACTAAAACAATTTTAATTTTGACGTTAGTT 234
 AAGTAATTTATATAGATGGATAAATTGAGCAAGCACATTACGAACCTGCGGATCAAGGAGTCAACAATTTTAATTCCTTA 312
 GGCCTCTAGACGAACCTGCGGAGCAAGG -> DZ2BGENF
 XbaI
 CGTTATACACAAAATTATCTAAATACTATATATATATACAGCTGCATGCTACGATAATGATCAAAATGTTTATGTACTT 390
 TTCAGCGGAAAATTCCTTGTCGCCATACATTACTGTGTTAATGAATCATTTAAATATGTGAAGGAGGAAAAGAGTACAAAA 468
 GGAGTTTGTGTGAGGCATTTCCGACACACTGAAATGTGAATAATAAAGGAATTGCCGAATGATTTTCTAGTTGGT 546
 GAAGTGGGTGAAAATTGTATGTCCATTGCTTATAAACTATAAAATATAATATNTTNATATTATCACTNTGGACATTAG 624
 TNNGATAGACCCCTAGCTAAAATTTTTTAAAAATTATACATTTCATTTTCTTNAAGTACCACAACTTAATTAATCACAATCGGA 702
 TAAAAATTGTTTAAGAAACCATTAACAACTCAGCTTGTGGACTCTGAGAGAACTAAGAGCTAGACATACGGTTAGTAG 780
 TGTAGCCGCATTTTTTATGCTTAATTGTCTTAAGCATGACTTCTATGCTCTCTGATGATATTTATTTTTAATATCCTAG 858
 GACATATGGATTTGATAAAGATCTTATCAACCTTTCAACAAGACCATTAGCTCAACAAACAAAATACTGAAAGTATAT 936

6/27

7/27

| | |
|---|------|
| AAATCTGGTTACAGAAATTCCTTATGCGCAAAAATATCATATAATATATAGAAATTCGGTTATGATTAAGATGAATTTTAA | 1013 |
| ATTAATATATATTTTTCACCTTTTGTTCCTTATGTATTCCTTAGTATTTTGTTCACCATATTGACCGATTGGTGTCATATTA | 1092 |
| GTTTGGTAAGACAACACTCAGTTGCAACGATGCAGATTACATTTTCAGGAAGATTTCATGTAAGAAAGATATTTCGCTTTGT | 1170 |
| GGTGTGAAAAATATGCCCTCTTTTCACCTTTTTCCTCAACTATAAATTTTCGATCGATGTATCTACGTTCTTAACACAATTTCAC | 1248 |
| AATCTTCTTTTAGAATCCAAAATGTAAAGCCGCTTTCTAATCTCTTTCTCAGTATACATATGTAATATGTATGCATATA | 1326 |
| TTATTATTTCATAATAACACGAAACCCATGCATGCAAGAAAGATAGTTACACGCTCATACAAACACAAAAAACATA | 1404 |
| CGCATGCATTAGAACACTTGTATGTAAATTTCCATAAATGTTTTCGATATAAACATTCCTTCGTTTAAATTAGCTTCCTTTT | 1482 |

| | |
|---|------------------|
| NcoI | |
| <- GTTCGAAGAAAAAACCAGCATGGCAG | DZ2BGENR AACCAAG |
| GTGTGAAGATTGTTCGAAGAAAAAACCAGAGATGGCAACAACGTCAACATCCACGGGAGATATCAAGAAACCAAG | 1560 |
| M A T T S T S T S T G D I K K T K | |

| |
|--|
| TCAGTAGAAGTG -> DZ2BFL |
| TCAGTAGAAGTGAAGAAGAACTTAACGTGTGTGATCGTTGTATGATGATACAGTAATTCGTAACTTCACGAGAATATC |
| S V E V K K L N V L I V D D D T V I R K L H E N I |
| ATCAAATCGATCGGTGGAATTTTCACAGACGGCTAAGAACCGGTGAGGAGGCAGTGAACATCCACCGCGACGGCAATGCA |
| I K S I G G I S Q T A K N G E E A V N I H R D G N A -> |

FIG. 5 CONT'D

Fig. 6

TCGTCNAT

GGCACGACGACGAATCGAAGATGGCAACAAATCCATGGGAGATATCGAGAAAATAAAGAAGAACTAAACGTGTGATCGTC

GATGATCCIGTAACGTAAACTTCACGAGATTATCATCAATCAATCGGTGA---ATTTCACAGACAGCTAAGAACGGTG
..... | | | | | . | | | | | . | | | | | . | | | | |
GATGATGCCACTAAACCTTATAATTCATGAGAAGATCATCAAAGCGATTGGGGGTATTTTCACAGACAGCGAATAACGGTG

90 | 100 | 110 | 120 | 130 | 140 | 150 | 160
AGGAGGCAGTGAACATCCACCGGACGGCAATGCATCTTTTCGACCCTTATCCTAATGGATAAAGAAATGCCCGAGAGGGATGG
.....|.....|.....|.....|.....|.....|.....|.....
AGGAGGCAGTAATCATCCACCGTGACGGCGGTCACTTTTGACCTTATCCTAATGGATAAAGAAATGCCCGAGAGGGATGG
| 170 | 180 | 190 | 200 | 210 | 220 | 230 | 240

ACTTTCGGCAACTAAGAAGCTAAGAGAAATGAAAGTGACGTCTATGATTATTGCGGTGACGCACACTGGCTGACAATGAAGAG
TGTTCGACAACTAAGAAGCTAAGAGAAATGGAAGTGAAGTCAATGATTGTTCGGGTGACTTCACTGGCTGACAATGAAGAG

9/27





FIG. 6 CONT'D



10 / 27






FIG. 6 CONT'D

11/27

FIG. 7

NORTHERN ANALYSIS OF EXPRESSION OF
DZ2B IN PODS AND OTHER TISSUES



DZ = POD DEHISCENCE ZONE (20 - 06 daa)
 NZ = POD NON-ZONE
 AZ = ABSCISSION ZONE
 F = FLOWER
 L = LEAF
 R = ROOT
 S = SEED

12 / 27




Fig. 8

FIG. 9

ATATATGTGATACAGATACATCTATATACAAATTAAACACGAAACCATACATGCACGGTGTGATCACACACGCACACA 78

CATAGAAACATAAACACGCAATAATTCTCTATACAGTTTAATTCATTTTAACTTACTTCTTTTTTTTGGTGAAGAT 156

TCTTGAGAGAAAGAAATCGAAGATGCGCAACAAAATCCACCGAGGTACCGAGAAAACCAAGTCGATAGAAAGTGAAGA 234
 M A T K S T G G T E K T K S I E V K K

AGAAACTAATCAACGTGTTGATCGTCGATGATCCATTAAACCGTAGACTCCACGAGATGATCATCAAAACGATCG 312
 K L I N V L I V D D D P L N R R L H E M I I K T I G

<- CGATCG

<- CGAATAACGGTGAGGAGGCATTAATCA GW1
 GAGGAATTTCAGACTGCAA AT3GW2
 GAGGAATTTCAGACTGCAAAAGAATGCGGAAGAGGCNGTGATCCTCCACCGTGACGGCGAAGCATCTTTCGACCTTA 390
 G I S Q T A K N G E E X V I L H R D G E A S F D L I

TTCTAATGGATAAGGAAATGCCCTGAGAGGGATGGAGTTTCGACAAATTAAGANGCTAAGAGAAATGAAAGGACGTCAA 468
L M D K E M P E R D G V S T I K X L R E M K G T S M

TGATCGTTGGGTAACGTACGTAGCTGACCAAGAAGAAGCGTAAGGCTTTTATGGAAGCTGGGCTCAACCATTCCT 546
I V G V T S V A D Q E E E R K A F M E A G L N H C L

TGGAAAAACCCCTTAACCAAGGCCAAGATCTTCCCGCTCATTTAGCCACCTCTTCGATGCTTGATGGAAGGCTCATT 624
E K P L T K A K I F P L I S H L F D A .

AATGTATCTATATTTTCAATCATGAAATCACCTACACGTGTATTTGACACAAAATCTGCATTGTTGTGATATAGGG 702

TTTCTCATATCTATGTTTGATTATTTTCTTATCGTCCGAGGTAAATCATGCAAGTCATTCTTTTGGCTAATAAAA 780

TATTAAATAAGGTTTCTCAAAAAAAAAAAAAAAAAA 818

FIG. 9 CONT'D

15 / 27

A) DZ2A13 peptide sequence (top)
DZ2 peptide sequence (bottom)

FIG. 10

| | | | | | | | |
|---|---|----|----|----|----|----|----|
| 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 |
| | | | | | | | |
| MATKSTGGTEKTSIEVKKKLINVLIVDDDDPLNRRLLHEMI | IKTIGGISQTAKNGEEXVILHRDGEASFDLILMDKEMPERDG | | | | | | |
| . | .. | .. | .. | .. | .. | .. | .. |
| MATKSMGDIKIKKKLNVLIVDDDDPLNLI | HEKIIKAIGGISQTANNNGEEAVIIHRDGGSSFDLILMDKEMPERDG | | | | | | |
| | | | | | | | |
| 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 |

| | | | | | |
|----------------------|--|-----|-----|-----|-----|
| 90 | 100 | 110 | 120 | 130 | 140 |
| | | | | | |
| VSTIKXLREMKGTSMIVGVT | SVADQEEERKAFMEAGLNHCLEKPLTKAKIFPLISHLFDA | | | | |
| ... | ... | ... | ... | ... | ... |
| VSTTKKLREMEVKSMIVGVT | SLADNEEERRAFMEAGLNHCLEKPLTKDKIIPLINQLMDA | | | | |
| | | | | | |
| 80 | 90 | 100 | 110 | 120 | 130 |

16 / 27

B) DZ2AT3 peptide sequence (top)
DZ2B peptide sequence (bottom)

| | | | | | | | |
|---|-----------------------------------|-----------------------------------|-----------|-----------|-----------|-----------|-----------|
| 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 |
| | | | | | | | |
| MATKSTGGTEKTSIEVKKKLINVLIVDDPLNRRRLHEMI | IKTIGGISQ | TAKNGEEXVILHRDGEASFDLILMDKEMPERDG | | | | | |
| | | | | | | | |
| VXDDPVTIRKLHEIIKSIGGISQ | TAKNGEAAVNIHRDGNASFDLILMDKEMPERDG | | | | | | |
| | 10 | 20 | 30 | 40 | 50 | | |

| | | | | | |
|----------------------|--|-----------|-----------|-----------|-----------|
| 90 | 100 | 110 | 120 | 130 | 140 |
| | | | | | |
| VSTIKXLREMKGTSMIVGVT | SVADQEEERKAFMEAGLNHCLEKPLTKAKIFPLISHLFDA | | | | |
| . | | | | | |
| LSATKKLREMKVTSMI | IGVTTLADNEEERKAFMEAGLNHCLEKPLSKAKILPLINNLMDA | | | | |
| | 60 | 70 | 80 | 90 | 100 |

FIG. 10CONT'D

FIG. 11

GTAATGCGACTCACTATAGGCGACGCGTGGTCGACGCCCGGCTGGTCCCTCATTCGTATMGGCCCAATGGGCTACT 78

CACTAGTAGGCGACGCGTGGTCG -> ATDZ2F

SpeI

AAAACAGTTTCACGATTGTTTTTTTTTTTTTTTAAATTTTAAACATGTA GTGGATAATTGGCTATAAATTATG 156

TAAAAAATTTACGATAGATTGTTGAATTTTCGAGTTAAATAATCTTCAAATTACCTCACATTTACAAAAA 234

GGTAGAACTGTTGAAAAACTAATGCTCTATAAAACACTAGACAATAACAAAAACGTAATGCGTAAAGAACCTAAATT 312

ATGATTTTATTTATCTTTCTTCTTTTCCGCGAGTATAAGCCATTTTTCATAGTAAAGCATTACGAAATACGACATTG 390

AACACTACTGACATATAAAGTAGTAGATTTTGATGGGTTAACTTGTA TGCTTAATTMGCTTAAGCATGAAC TTCAATG 468

CTTTTATAAAAGTACTTTCATGAGAAATATTCCTCGTTCTATACTAGCAGAAGGTT CGATAGTGATTTTACAA CCGTTC 546

AACAAAACCTTTAAACCCAAAACCAAGAAATGAAAGTATCTAAACTTGATTA TACATTTCTTGCTAAATTATCAA 624

ATAACATACTCTCTTTTGTTTACTTATAAACGATATGAAAGAAATAAATAAAAAGAACATAGAATCTTATTATGATCT 702

18/27

AGAAGAATTAAAGAAATATATATATATTTTTTTCATTTCCTACTCATGTTCTTATACATTCTTTAAATTGTT 780
 CACCATTTTGATTACTTGTTCTCATATTAGTTTGTTATACAACCTCACTTAGAATAATGTAGATTACATTTTCAGCCAA 858
 ATTCATGTAAAGATGCTTTTCTTTGTGATGTTTAAATGCTTTCTTTTCACCTTTTTCCTTTTAACTATAAAT 936
 CTTGATCGAATGCCCTACCTTCTTAGAACATAAGATCTTCTTTTAAATCCAAAATCGTAGGCCACTATTTTCATTATACT 1013
 TATGTAATATATGTGATACAGATACATNTATATACAAATTAAACACGAAACCATACATGCACGGTGTGATCACACACG 1092
 CACACACATAGAAACATAAACACGCAATAATTCTATACAGTTTAATTTCATTTTAACTTACTTCTTTTCTTTTGGT 1170

NcoI

<- CTTGAGAGAAAGAAATCGACCATGGA
 GAAGATTCTTGAGAGAAAGAAATCGAAGATGGCAACAAAATCCACCGGAGGTACCGAGAAACCAAGTCGATAGAAG 1248
 M A T K S T G G T E K T K S I E V
 TGAAGAAGAACTAATCAACGTGTTGATCGTCGATGATCCATTAAACCGTAGACTCCACGAGTGTTCATCAAAA 1324
 K K K L I N V L I V D D D P L N R R L H E C H Q ->

FIG. 11 CONT'D

19 / 27




FIG.12

20 / 27




FIG.13A

21 / 27




FIG.13B

22 / 27




FIG.14

FIG. 15

GGCATCAGGAGGTACCCGTAATCCCACCATACAACAAGTTCTGTGAAGTCTCCCA 60
AACTGCCAAGAGTCTCATATTAGTTCTTACTCTCAGAAATAAACACAGTGTCTCTGAA 120
AAGATTAGCGTTTCAAAACCCGAAATGGCCCGTTGTCAATGGAAGTCTTGCTATTTCTTA 180
M A R C H G S L A I F L 12
TGCGTTCTTTTGATGCTCGCTTGCTGCCAAGCTTTGAGTAGCAACGCTAGATGATGGATAT 240
C V L L M L A C C Q A L S S N V D D G Y 32
GGTCATGAAGATGGAAGCTTCGAAACCGATAGTTTAATCAAGCTCAACAACGACGAC 300
G H E D G S F E T D S L I K L N N D D 52
GTTCTTACCTTGAAAAGCTCCGATAGACCCACTACCGAATCATCAACTGTTAGTGTTCG 360
V L T L K S S D R P T T E S S T V S V S 72
AACTTCGGAGCAAAAGGTGATGGAAAAACCGATGATACCTCAGGCTTTCAAGAAAGCATGG 420
N F G A K G D G K T D D T Q A F K K A W 92
AAGAAGGCATGTTCAACAAATGGAGTGACTACTTTCTTGATTCCTAAAGGGAAGACTTAT 480
K K A C S T N G V T T F L I P K G K T Y 112
CTCCTTAAGTCTATTAGATTCAGAGGCCCATGCAAAATCATACGTAGCTTCCAGATCCTA 540
L L K S I R F R G P C K S L R S F Q I L 132
GGCACTTTATCAGCTTCTACAAAACGATCGGATTACAGTAATGACAAGAACCCTGGCTT 600
G T L S A S T K R S D Y S N D K N H W L 152

23/27

24 / 27

ATTTGGAGACGCTTAATCTATCAATCGATGGCGCTCGCGGGGATTGTTGATGGC 660
 I L E D V N N L S I D G G S A G I V D G 172

AACGGAAATCTGGTGGCAAACTCATGCAAAATCGACAAATCTAAGCCATGCACAAA 720
 N G K I W Q N S C K I D K S K P C T K 192

GCGCCAACGGCTCTACTCTCTACAACCTAAACAATTGGAATGTGAAGAAATCTGAGAGTG 780
 A P T A L T L Y N L N N L N V K N L R V 212

AGAAATGCACAGCAGATTTCGATTTGAGAAATGCAACAGTGTGTGATGTTAAGAAT 840
 R N A Q Q I I S I E K C N S V D V K N 232

GTTAAGATCACTGCTCGCGGATAGTCCCAACACGGATGGTATTCATATCGTTGCTACT 900
 V K I T A P G D S P N T D G I H I V A T 252

AAAACATTGCAATCTCCAATTCAGACATTGGGACAGGTGATGATTCATATCCATTGAG 960
 K N I R I S N S D I G T G D D C I S I E 272

GATGGATCGCAAAATGTTCAAAATCAATGATTAACTTGGCGGCCCGGTGATGGCATCAGC 1020
 D G S Q N V Q I N D L T C G P G H G I S 292

ATTGGAAGCTTGGGGGATGACAAATCCAAAGCTTATGTATCGGGAATTAATGTGGATGGT 1080
 I G S L G D D N S K A Y V S G I N V D G 312

FIG. 15 CONT'D

25 / 27

GCTACGCTCTCTGAGACTGACAAATGGAGTAAGAAATCAAGACTTACCAGGGAGGTCAGGA 1140
 A T L S E T D N G V R I K T Y Q G G S G 332

ACTGCTAAGAACATTAATAATCCAAACATTCGTATGGATAATGTCAAGAAATCCGATCATA 1200
 T A K N I K F Q N I R M D N V K N P I I 352

ATCGACCAGAACTACTGCGACAAGGACAAAATGCGAACAACAAGAAATCTGCGGTTCAAGTG 1260
 I D Q N Y C D K D K C E Q Q E S A V Q V 372

AACAAATGTCGTCTATCGGAACATACAAGGTACGAGCGCAACGGATGTGGCGATAATGTTT 1320
 N N V V Y R N I Q G T S A T D V A I M F 392

AATTGCAGTGTAATAATCCATGCCAAGGTATTGTGCTTGAGAAATGTGAACATCAAAGGA 1380
 N C S V K Y P C Q G I V L E N V N I K G 412

GGAAAAGCTTCTTGCAAAAATGTCAATGTTAAGGATAAAGGCACCGTTTCTCCTAAATGC 1440
 G K A S C K N V N V K D K G T V S P K C 432

CCTTAATTACTAAGTTGATTATGTAATATACATAAAATACGTATTATATGTGTTATAGAT 1500
 P 433

GCCATCTATATCCTTATCTACGTATTGATTCTCGATATATAGAAACTAAGGATTAT 1560

GGGAAATATACATAACAATAGTTGAGATAATTGTTGCTTGTATATGGTTCACCTGAAGTTGA 1620

TTGCTTGTCCACGAATAAATGAATAATGTCATTGTC 1657

FIG. 15 CONT'D

26 / 27

aggtgaccggttgctgatggcaatgtgctgggtcaagcgaagaggtagacgggtggcttgagaga
V T V A D G N V L V K R E V D G G L E T
cagttaaagtcaaattgccagctgtcatttagcgccgacttgcggtcgaatgagccgcgggt
V K V K L P A V I S A D L R L N E P R Y
acgctactctgccccaatatcatgaaggccaagaagaagcccatcaaaaagctcacagcca
A T L P N I M K A K K K P I K K L T A T
cagatgtcgggtgtggacttggcgccacgtcaacaagtgttgagcgtagaagacccgcccc
D V G V D L A P R Q Q V L S V E D P P T
ccagacaggctgggttccattgtgcctgatgtcgacactctcatcaccaagttgaaagaaa
R Q A G S I V P D V D T L I T K L K E K
agggtcatttgtaatgcaatgtcaccaatacagttgttttagttcttaciaaattcttcgt
G H L *
gaggttttcagctgttaccaataatatTTTTTcaaaatcgattttattttacttgtaatt
taaaagatcaaataattaatacaatgaacatttttgtaacagcaatcttttttttatattt
tggagatttcacgcacttatgtcataattatTTTTtatcaatttattgttgtttgtagtg
atataataaagtatgttttctgggtcaaaa

FIG. 16

27/27

OSR79 306bp

```

5'  gggtgggtcgaaccataggtggaaagcttcttctctctcgcttgacaaatcctctgggtt
    L G R T I G G K L L S L S L D K S S G S
      . . . . .
cgggttttcagtcgccatcaggagtttctctctatgggtaaagctgaggttcaaatgaaacttg
    G F O S H Q E F L Y G K A E V O M K L V
      . . . . .
tccctggtaactctgtggaaacagtcacaacattctctttaaatacccggaactacat
    P G N S A G T V T T F Y L K S P G T T W
      . . . . .
gggatgagatcgatttcgagttcttgggaaacataaagtggccatccctatactctccata
    D E I D F E F L G N I S G H P Y T L H T
      . . . . .
ctaattgtttacacacgaaggctctggagacaaagaacagcagtttcatctatgggttcgac
    N V Y T R R L W R Q R T A V S S M V R P
      . . . . .
ccgacc      3'
      D

```

FIG. 17

INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/GB 99/00905

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/29 C12N15/82 C12N15/11 C12N5/10 C07K14/415
A01H5/00 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|----------------------------|
| X | NELSON D E ET AL: "Abundant accumulation of the calcium-binding molecular chaperone calreticulin in specific floral tissues of Arabidopsis thaliana." PLANT PHYSIOLOGY, (1997 MAY) 114 (1) 29-37. , XP002111979 see esp. p.36 2.par. - end; abstract --- | 1-3, 10-17, 25,26,29 |
| A | WO 97 13865 A (PRINSEN ELS ;BOTTERMAN JOHAN (BE); ONCKELEN HENRI VAN (BE); PLANT) 17 April 1997 (1997-04-17) the whole document --- | 1-29 |
| A | WO 96 30529 A (JENKINS ELIZABETH SARAH ;ROBERTS JEREMY ALAN (GB); NICKERSON BIOCE) 3 October 1996 (1996-10-03) cited in the application the whole document --- | 1-29 |
| -/-- | | |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

17 August 1999

Date of mailing of the international search report

30/08/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 99/00905

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|-----------------------|
| A | WO 94 23043 A (COUPE SIMON ALLAN ;ROBERTS JEREMY ALAN (GB); ISAAC PETER GEOFFREY) 13 October 1994 (1994-10-13) cited in the application the whole document ---- | 1-29 |
| P,X | WHITELAW C A ET AL: "An mRNA encoding a response regulator protein from Brassica napus is up-regulated during pod development" JOURNAL OF EXPERIMENTAL BOTANY, (MAR 1999) VOL. 50, NO. 332, PP. 335-341., XP002111980 the whole document ----- | 1-29 |

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 99/00905

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9713865 A | 17-04-1997 | AU 7284796 A | 30-04-1997 |
| | | CA 2234222 A | 17-04-1997 |
| | | CN 1211282 A | 17-03-1999 |
| | | CZ 9801042 A | 16-09-1998 |
| | | EP 0853676 A | 22-07-1998 |
| | | HU 9802535 A | 01-02-1999 |
| | | PL 326082 A | 17-08-1998 |
| WO 9630529 A | 03-10-1996 | AU 5155996 A | 16-10-1996 |
| | | CA 2216805 A | 03-10-1996 |
| | | EP 0817857 A | 14-01-1998 |
| WO 9423043 A | 13-10-1994 | AU 6381994 A | 24-10-1994 |
| | | CA 2159614 A | 13-10-1994 |
| | | EP 0692030 A | 17-01-1996 |
| | | US 5907081 A | 25-05-1999 |